



Eur päisches Patentamt
Eur pean Patent Office
Offic eur péen des brevets

Publicati n number:

0 187 433
A1

EUROPEAN PATENT APPLICATION

Application number: 85300205.3

Date of filing: 11.01.85

Int. CL⁴: **A 61 K 9/06**
A 61 K 47/00

Date of publication of application:
16.07.86 Bulletin 86/29

Designated Contracting States:
CH DE FR GB IT LI NL

Applicant: **TEIJIN LIMITED**
11 Minami Honmachi 1-chome Higashi-ku
Osaka-shi Osaka-fu(JP)

Inventor: **Suzuki, Yoshiki**
5-20-2, Tamadaira
Hino-shi Tokyo(JP)

Inventor: **Ikura, Hiroshi**
Gurin-kopu Hino 2-501,2-14-1
Shinmei Hino-shi Tokyo(JP)

Representative: **Arthur, Bryan Edward et al,**
Withers & Rogers 4 Dyer's Buildings Holborn
London EC1N 2JT(GB)

Powdery pharmaceutical composition suitable for application to mucosa of oral or nasal cavity.

A powdery pharmaceutical composition for application to the mucosa of the oral or nasal cavity including: (a) a base selected from the group consisting of lower alkyl ethers of cellulose; (b) a pharmaceutically effective amount of a drug selected from the group consisting of steroid or glycyrrhizic acid type anti-inflammatory agents; and (c) a stabilizer for the drug composed of at least one low irritant solid organic acid selected from the group consisting of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, unsaturated lower aliphatic monocarboxylic acids having six or less carbon atoms, and cellulose derivatives having carboxyl group.

This powdery pharmaceutical composition has extremely excellent stability in the steroid or glycyrrhizic acid type anti-inflammatory agent contained therein, low irritation of the mucosa of the oral or nasal cavity, and sustained release of the drug over an extended period of time.

EP 0 187 433 A1

POWDERY PHARMACEUTICAL COMPOSITION SUITABLE FOR
APPLICATION TO MUCOSA OF ORAL OR NASAL CAVITY

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a powdery pharmaceutical composition suitable for application to the mucosa of the oral or nasal cavity. More specifically, it relates to a powdery pharmaceutical composition, suitable for application to the mucosa of the oral or nasal cavity, containing (a) lower alkyl ethers of cellulose as a base, (b) a steroid or glycyrrhizic acid type anti-inflammatory agent as a drug component, and (c) a solid low irritant organic acid as a stabilizer for the drug component.

The powdery pharmaceutical composition exhibits extremely excellent stability in the steroid or glycyrrhizic acid type anti-inflammatory agent contained therein, low irritation of the mucosa of the oral or nasal cavity, and sustained release of the drug over an extended period of time. Accordingly, the powdery pharmaceutical composition according to the present invention can be effectively used for the treatment of oral diseases such as stomatitis, erosion or sores generated by radiotherapy, and lichen planus or nasal diseases such as allergic rhinitis and vasomotor rhinitis.

25 2. Description of the Related Art

Various pharmaceutical preparations, such as buccal tablets, troche tablets, sublingual tablets, and oral ointments, have been heretofore known in the art for application to the oral cavity. Buccal tablets, troche tablets, and sublingual tablets, however, give a foreign feeling to the oral cavity. Therefore, patients often chew on and swallow the tablets. Oral ointments, on the other hand, usually contain mixtures of beeswax

or plastibase with gelatin, pectin, or sodium carboxymethyl cellulose. These mixtures are unsatisfactory because the adhesion properties thereof to the oral mucosa are not sufficient and also because they have
5 unpleasant tastes.

On the other hand, various pharmaceutical preparations, such as nasal ointments, jellies, nose drops, and sprays, have been known in the art for application to the nasal cavity. Nasal ointments and
10 jellies, however, are difficult to apply to deep parts of the nasal cavity, such as the concha nasalis superior. With, nose drops and sprays, the active components do not remain in the nasal cavity for an extended period of time.

Various sustained release pharmaceutical preparations for oral or nasal cavity administration are known in the art. With these, the efficacy of the drugs is spread over a long period of time by the gradual or sustained releasing the active drug components from the
15 pharmaceutical preparations. For example, Japanese Unexamined Patent Publication (Kokai) No. 57-118511 discloses powdery sustained release preparations suitable for adhesion to the oral cavity. These preparations use, as a base, hydroxypropyl cellulose and have the
20 advantages of an excellent sustained release of the drugs and possible application over a wide range of the oral cavity. Furthermore, U.S. Patent No. 4,294,829 discloses powdery sustained release preparations suitable for application to the nasal cavity. These contain lower
25 alkyl ethers of cellulose and drugs and also have the advantage of excellent sustained release of the drugs.

It is, however, more desirable to improve the stability of drugs in the above-mentioned sustained release preparations for application to the mucosa of
30 the oral or nasal cavity.

SUMMARY OF THE INVENTION

Accordingly, the object of the present invention is

to provide a powdery pharmaceutical composition, suitable for application to the mucosa of the oral or nasal cavity, having extremely excellent stability of the carried drug, i.e., a steroid or glycyrrhizic acid type anti-inflammatory agent.

Another object of the present invention is to provide a powdery pharmaceutical composition, suitable for application to the mucosa of the oral or nasal cavity, containing a lower alkyl ether of cellulose as a base, a steroid or glycyrrhizic acid type anti-inflammatory agent as a drug, and a specific solid low irritant organic acid as a stabilizer for the drug.

A further object of the present invention is to provide a powdery pharmaceutical composition having low irritation of the mucosa of the oral or nasal cavity.

A still further object of the present invention is to provide a powdery pharmaceutical composition having excellent sustained release of the drug.

A still further object of the present invention is to provide a powdery pharmaceutical composition having a particle diameter suitable for administration to the oral or nasal cavity.

A still further object of the present invention is to provide a method for preparing a powdery pharmaceutical composition suitable for application to the mucosa of the oral or nasal cavity.

A still further objects of the present invention is to provide a method for stabilizing a powdery pharmaceutical composition containing a steroid or glycyrrhizic acid type anti-inflammatory agent as a drug.

Other objects and advantages of the present invention will be apparent from the following description.

According to the present invention, the above-mentioned objects and advantages can be achieved by a powdery pharmaceutical composition including: (a) a base selected from the group consisting of lower alkyl ethers of cellulose; (b) a pharmaceutically effective

amount of a drug selected from the group consisting of steroid or glycyrrhizic acid type anti-inflammatory agents; and (c) a stabilizer for the drug composed of at least one low irritant solid organic acid selected from the group consisting of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, unsaturated lower aliphatic monocarboxylic acids having six or less carbon atoms, and cellulose derivatives having carboxyl groups.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

According to the present invention, the desired powdery pharmaceutical composition, i.e., a composition having extremely excellent stability of the drug, low irritation to the mucosa of the oral or nasal cavity, and excellent sustained release of the drug, can be obtained by using a lower alkyl ether of cellulose as a base, a steroid or glycyrrhizic acid type anti-inflammatory agent as a drug, and a specific solid low irritant organic acid as a stabilizer for the drug, i.e., by combining the specified base and the specified drug with the specified organic acid.

The lower alkyl ethers of cellulose usable as a base in the present invention are those obtained by at least partially substituting the same or different lower alkyl ether groups for a plurality of hydroxyl groups of cellulose. The lower alkyl groups in the lower alkyl ether groups may be substituted with substituents. An example of such preferable substituents is a hydroxyl group.

Preferable examples of the optionally substituted lower alkyl groups are a methyl group, and hydroxy lower alkyl groups having 2 or 3 carbon atoms.

Examples of the optionally substituted lower alkyl groups are methyl, ethyl, n-propyl, iso-propyl, β -hydroxyethyl, and β -hydroxypropyl.

Examples of the optionally substituted lower alkyl ethers of cellulose are methyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, and hydroxypropylmethyl cellulose. Of these lower alkyl ethers of cellulose, the use of methyl cellulose, hydroxypropyl cellulose, and hydroxypropylmethyl cellulose is preferable due to the facts that these lower alkyl ethers of cellulose have substantially no odor and irritation when applied to the mucosa of the oral or nasal cavity, which is particularly sensitive to odor and irritation, and that the desired sustained release of the drug can be obtained. Of these lower alkyl ethers of cellulose, the use of hydroxypropyl cellulose is especially preferable.

The lower alkyl ethers of cellulose can be used alone or in any mixture thereof.

When the above-mentioned lower alkyl ethers of cellulose are used as a base in the powdery pharmaceutical composition according to the present invention, the pharmaceutical preparations thus obtained can adhere to the mucosa of the oral or nasal cavity and will swell to gradually release the active drug components therefrom over a long period of time. Thus, the affected or diseased parts can be treated by the drugs over a long period of time.

Although there is no limitation in the molecular weight of the lower alkyl ethers of cellulose usable in the present invention, lower alkyl ethers having a viscosity in 2% aqueous solution thereof of 3 to 10,000 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$, more preferably, 1000 to 4000 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$, are preferably used.

In preparing unit dosage forms of powdery pharmaceutical compositions in which lower alkyl ethers of cellulose are contained as a base, the compositions are generally filled in hard capsules. These capsules are set in a specially designed spraying device. The spraying device is provided with needle for spraying the powdery pharmaceutical compositions into the oral or

nasal cavity. In this case, when the moisture content of the lower alkyl ethers of cellulose in the capsules is too small, the moisture contained in the membranes of the capsules tends to be absorbed by the lower alkyl ethers of cellulose and, as a result, the capsules become too rigid. Therefore, lower alkyl ethers of cellulose having a moisture content of 3% to 9% by weight, especially 4% to 8% by weight, are preferably used as a base.

10 The drugs usable in the present powdery pharmaceutical compositions are steroid or glycyrrhizic acid type anti-inflammatory agents. Examples of such steroid type anti-inflammatory agents are beclomethasone dipropionate, betamethasone, betamethasone valerate, 15 triamcinolone, triamcinolone acetonide, dexamethasone, fluocinolone acetonide, fluocinonide, flumethasone, hydrocortisone, prednisolone, and prednisone. Examples of glycyrrhizic acid type anti-inflammatory agents are glycyrrhizic acid, disodium glycyrrhizinate, trisodium 20 glycyrrhizinate, monopotassium glycyrrhizinate, dipotassium glycyrrhizinate and monoammonium glycyrrhizinate. These drugs can be used alone or in any mixture thereof in the present powdery pharmaceutical composition.

According to the present invention, the above-mentioned anti-inflammatory agents contained in the 25 powdery pharmaceutical compositions comprising the above-mentioned lower alkyl ethers of cellulose as a base can be advantageously stabilized by incorporating thereinto, as a stabilizer, a specified organic acid, 30 i.e., at least one solid low irritant organic acid selected from the group consisting of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, 35 unsaturated lower aliphatic monocarboxylic acids having six or less carbon atoms, and cellulose derivatives having carboxyl groups.

The organic acids usable as a stabilizer for the drug in the present invention are those selected from the above-mentioned group of the organic acids which are solid at room temperature, less irritant to the mucosa of the oral or nasal cavity, and substantially odorless or with only a very slight odor. By the term "organic acids" is meant acids comprising organic groups and free carboxylic groups. Use of the organic acids solid at room temperature facilitates the production of the desired powdery pharmaceutical compositions in a uniform particle size. Use of the less irritant organic acids means higher safety when the composition is applied to the mucosa of the oral or nasal cavity.

Examples of the preferable saturated higher aliphatic monocarboxylic acids are saturated higher aliphatic hydrocarbon monocarboxylic acid having 12 to 24 carbon atoms, such as stearic acid, palmitic acid, lauric acid, and myristic acid. Examples of the preferable aliphatic polycarboxylic acids are aliphatic hydrocarbon dicarboxylic acids having 4 to 6 carbon atoms, such as, succinic acid, fumaric acid, and maleic acid. Examples of the preferable hydroxy aliphatic polycarboxylic acid are hydroxy aliphatic hydrocarbon di- or tri-carboxylic acids having 4 to 8 carbon atoms, such as, tartaric acid and citric acid. Examples of the preferably aromatic acids are benzoic acid and its derivatives, such as parahydroxy benzoic acid and salicylic acid. Examples of the preferable unsaturated lower aliphatic monocarboxylic acids are unsaturated lower aliphatic hydrocarbon monocarboxylic acid having 4 to 6 carbon atoms, such as sorbic acid. Examples of the preferable cellulose derivatives having carboxyl groups are carboxymethyl cellulose and carboxymethylethyl cellulose. These organic acids can be used alone or in any mixture thereof.

Of the above-mentioned organic acids, saturated higher aliphatic monocarboxylic acids having 12 or more

carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, and cellulose derivatives having carboxyl groups are preferably used. For powdery pharmaceutical compositions for application to the nasal cavity, use of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms and cellulose derivatives having carboxyl groups is especially preferably from the viewpoints of low irritation and substantial odorlessness.

There is no specific limitation in the amount of the above-mentioned organic acids in the present pharmaceutical composition as long as the anti-inflammatory agents are stabilized. The preferable amount of the organic acid in the present pharmaceutical composition is within the range of from 0.05% to 5% by weight, especially from 0.1% to 3% by weight, based upon the total weight of the composition. Although there is also no critical limitation in the particle size of the organic acid, the organic acid preferably has such a particle size that at least about 90% by weight of the powder particles has an effective particle diameter of about 75 μm or less, more preferably about 20 μm to about 75 μm , to obtain a powdery pharmaceutical composition capable of being effectively administered to the mucosa of the oral or nasal cavity.

As mentioned above, the powdery pharmaceutical compositions according to the present invention contain, as essential constituents, the above-mentioned lower alkyl ethers of cellulose, steroid or glycyrrhizic acid type anti-inflammatory agents, and organic acid stabilizers. Although there is no specific limitation in the particle size of the present pharmaceutical composition, the particle size of the present pharmaceutical composition is preferably such that at least about 90% by weight of the powder particles thereof has an effective particle diameter of about 20 μm to about 250 μm . The present powdery pharmaceutical composition

having such a particle size distribution is preferably due to the fact that the amount of the powder particles adhering to the mucosa of the oral or nasal cavity becomes large. That is, when the pharmaceutical composition having the above-mentioned preferable particle size distribution is applied to the oral or nasal cavity, the amount of the pharmaceutical composition reaching, for example, the lung becomes small or the amount of the pharmaceutical composition scattered out of the oral or nasal cavity becomes small. Furthermore, the pharmaceutical composition having the above-mentioned preferable particle size can well adhere to the mucosa of the oral or nasal cavity and can be effectively and gradually released. The powdery pharmaceutical composition according to the present invention more preferably has such a particle size distribution that at least about 90% by weight of the powder particles has an effective particle diameter of about 20 μm to about 150 μm .

The term "effective particle diameter" used herein means that determined by the opening sizes of sieves. For example, a powder having an effective diameter (d) of $37 < d \leq 44$ passes through a sieve having an opening size of 44 microns but does not pass through a sieve having an opening size of 37 microns.

A vibratory sieve is used when the effective particle diameter of a powder is more than 37 microns, and a sonic sieve (Micro Hand Sifter SWM-2, a product of Tsutsui Rikagaku Kikai Co., Ltd.) is used when the effective particle diameter of a powder is not more than 37 microns.

The sieves used has an opening size of 500, 420, 350, 297, 250, 210, 177, 149, 125, 105, 88, 74, 63, 53, 44, 37, 25, and 20 microns, respectively.

The powdery pharmaceutical composition according to the present invention can optionally contain, in addition to the above-mentioned lower alkyl ethers of cellulose, anti-inflammatory agents, and organic acids, any conven-

tional ingredients for improving the physical properties, visual appearance, or odor of the pharmaceutical preparations. Examples of such ingredients are: lubricants such as talc and waxes; binders such as starch, dextrin, 5 tragacanth gum, gelatin, polyvinylpyrrolidone, and polyvinyl alcohol; diluents such as starch, crystalline cellulose, dextrin, lactose, mannitol, sorbitol, and anhydrous calcium phosphate; odor improvers such as menthol and citrus perfumes; preservatives such as 10 methyl p-hydroxybenzoate, propyl p-hydroxybenzoate; and various surfactants.

The powdery pharmaceutical composition according to the present invention can be prepared in any conventional manner. For example, the pharmaceutical composition can 15 be prepared by mechanically mixing the above-mentioned three essential constituents, i.e., the lower alkyl ethers of cellulose, the anti-inflammatory agents, and the solid organic acids and the above-mentioned optional ingredients, if any. The mixing of the composition can 20 be carried out by using any conventional means such as ball mills, and V-type mixers. The desired pharmaceutical composition can also be prepared by mechanically mixing all the above-mentioned constituents of the composition, compressing the resultant mixture under 25 pressure, e.g., 0.2 to 0.4 ton, to and pulverizing the compressed mixture so as to obtain the desired particle size distribution. The mixed or pulverized powder particles can be sieved to obtain the powdery composition having the desired particle size distribution, 30 if necessary.

The desired pharmaceutical composition can further be prepared by impregnating the lower alkyl ethers of cellulose with a solution or dispersion, in an organic solvent (e.g., methanol, ethanol, ethyl cellosolve, and 35 dichloromethane), of the anti-inflammatory agents and the organic acids and, if any, the optional ingredients, followed by drying the impregnated product. The said

organic solvent may contain water. The impregnation can be carried out by, for example, spraying the above-mentioned solution or dispersion to the lower alkyl ethers of cellulose in the form of powder particles
5 in a spray apparatus. The particle size distribution of the lower alkyl ethers of cellulose can be controlled to the desired range prior to the impregnation. Alternatively, the powder particles of the impregnated product can be pulverized and/or sieved after the impregnation
10 so as to obtain the desired particle size distribution.

Furthermore, the desired pharmaceutical compositions can be prepared by dissolving or dispersing the lower alkyl ethers of cellulose, the anti-inflammatory agents, the solid organic acid, and any optional ingredients
15 in organic solvents such as methanol, ethanol, ethyl cellosolve, and dichloromethane, water, or the mixture thereof followed by evaporating the organic solvents off. The resultant composition can be pulverized and/or sieved to obtain the desired particle size distribution.

20 According to the present invention, a powdery preparation in a unit dosage form including the above-mentioned powdery pharmaceutical composition can be obtained. The present pharmaceutical composition can be directly converted to the powdery preparation in the form of a unit dosage. The unit dosage of the powdery
25 preparations according to the present invention is preferably about 10 mg to about 400 mg, especially about 30 mg to about 300 mg, when applied to the oral cavity and about 5 mg to about 200 mg, especially about 10 mg
30 to about 100 mg, when applied to the nasal cavity.

The amount of the steroid or glycyrrhizic acid type anti-inflammatory agent in the powdery preparation in the unit dosage form depends upon the pharmaceutically effective amount of the anti-inflammatory agent to be
35 used and the number of dosages of the preparation. For example, when beclomethasone dipropionate is incorporated into the powdery preparations for application to the

nasal or oral cavity, the appropriate amount is about 25 to 200 μ g/day, although the conventional amount is about 200 to 400 μ g/day. When dipotassium glycyrrhizinate is incorporated into the powdery preparations for application to the oral cavity, the appropriate amount is about 0.1 to 15 mg/day, although the conventional amount is about 10 to 30 mg/day. This is because, since lower alkyl ethers of cellulose are used in the present pharmaceutical composition, a desired sustained release of the drug component can be obtained.

The powdery preparations of the present invention can be preferably filled into capsules, such as hard gelatin capsules, as a preferred dosage form.

The present powdery preparations can be advantageously applied to the oral or nasal cavity. For example, a capsule filled with the present powdery preparation is set in a spraying device equipped with a needle. The capsule is pierced with the needle to provide minute holes on the top and bottom sides. The powdery preparation is then sprayed or jetted out from the capsule by sending air into the capsule by means of, for example, a rubber ball.

As mentioned above, when the moisture content in the powdery preparation filled in a hard gelatin capsule is too small, the powdery preparation absorbs moisture from the gelatine membrane. Therefore, water is preferably contained in an amount of 3% to 8.5% by weight, especially 4% to 8.5% by weight, in the lower alkyl ethers of cellulose in the powdery preparations.

The powdery preparations according to the present invention can be widely used for the treatment of oral diseases such as stomatitis, erosion or sores generated by radiotherapy, and lichen planus or nasal diseases such as allergic rhinitis and vasomotor rhinitis.

EXAMPLE

The present invention will now be described in detail with reference to, but is by no means limited to,

the following examples, wherein all percentages and parts are on the weight basis unless otherwise specified.

Example 1

5 A 100 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 1550 cps at 37°C \pm 0.2°C and having a particle size such that 90% or more of the powder particles had a particle diameter of 37 to 149 μ m was used.

10 One part of the hydroxypropyl cellulose and 0.025 parts of beclomethasone dipropionate were thoroughly mixed in a mortar and, then, 9 parts of hydroxypropyl cellulose was gradually added thereto. The resultant mixture was thoroughly mixed and, then, the remaining hydroxypropyl cellulose were added and thoroughly mixed
15 in a mixer. Thereafter, 0.5 parts of stearic acid was added and thoroughly mixed.

Thus, a powdery composition for the application to the oral cavity containing beclomethasone dipropionate uniformly dispersed therein was prepared. The powdery
20 composition was filled in #2 hard gelatin capsules in an amount of 200 mg each to prepare a powdery preparation in a unit dosage form for the application to the oral cavity.

Example 2

25 A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Example 1, except that palmitic acid was used instead of stearic acid.

Example 3

30 A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Example 1, except that citric acid was used instead of stearic acid.

Example 4

35 A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Example 1, except that benzoic acid was

used instead of stearic acid.

Example 5

A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Example 1, except that sorbic acid was used instead of stearic acid.

Example 6

Powdery compositions for the application to the oral cavity were prepared in the same manner as in Example 1, except that the amounts of stearic acid were changed to 0, 0.1, 0.3, 0.5, and 0.7 parts. These powdery compositions were filed in #2 hard gelatin capsules in an amount of 200 mg each.

The capsules thus prepared were allowed to stand at a temperature of 60°C for one month. The residual percentages of beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 1.

Table 1

Content of stearic acid (%)	0	0.1	0.3	0.5	0.7
Residual percentage of beclomethasone dipropionate (%)	75.0	89.0	95.2	97.8	99.2

Example 7

Powdery compositions for the application to the oral cavity were prepared in the same manner as in Example 2, except that the amounts of palmitic acid were changed to 0, 0.1, 0.3, and 0.5 parts. These powdery compositions were filled in #2 hard gelatin capsules in an amount of 200 mg each.

The capsules thus prepared were allowed to stand at a temperature of 60°C for one month. The residual percentages of beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 2.

Table 2

Content of palmitic acid (%)	0	0.1	0.3	0.5
Residual percentage of beclomethasone dipropionate (%)	78.5	90.0	95.2	98.1

10 Example 8

Powdery compositions for the application to the oral cavity were prepared in the same manner as in Example 3, except that the amounts of citric acid were changed to 0, 0.05, 0.1, 0.5, 1, and 3 parts. These
15 powdery compositions were filled in #2 hard gelatin capsules in an amount of 200 mg each.

The capsules thus prepared were allowed to stand at a temperature of 60°C for one month. The residual percentages of beclomethasone dipropionate in the
20 capsules were determined.

The results were as shown in Table 3.

Table 3

Content of citric acid (%)	0	0.05	0.1	0.5	1	3
Residual percentage of beclomethasone dipropionate (%)	78.5	92.0	94.3	100.0	100.0	100.0

Example 9

Powdery compositions for the application to the oral cavity were prepared in the same manner as in Example 4, except that the amounts of benzoic acid were
35 changed to 0, 0.05, 0.1, and 0.5 parts. These powdery compositions were filled in #2 hard gelatin capsules in an amount of 200 mg each.

Th capsules thus prepared w re allowed to stand at a temperature of 60°C for one month. The residual percentages of beclomethasone dipropionate in the capsules were determined.

5 The results were as shown in Table 4.

Table 4

Content of benzoic acid (%)	0	0.05	0.1	0.5
Residual percentage of beclomethasone dipropionate (%)	82.5	93.2	94.0	100.0

Example 10

15 Powdery compositions for the application to the oral cavity were prepared in the same manner as in Example 5, except that the amounts of sorbic acid were changed to 0, 0.1, 0.3, and 0.5 parts. These powdery compositions were filled in #2 hard gelatin capsules in
20 an amount of 200 mg each.

The capsules thus prepared were allowed to stand at a temperature of 60°C for one month. The residual percentages of beclomethasone dipropionate in the capsules were determined.

25 The results were as shown in Table 5.

Table 5

Content of sorbic acid (%)	0	0.1	0.3	0.5
Residual percentage of beclomethasone dipropionate (%)	83.1	92.6	98.2	98.9

Example 11

35 Powdery preparations for the application to the oral cavity were prepared in the same manner, except that 100 parts of methyl cellulose having a viscosity of

a 2% aqueous solution thereof of 1335 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$ and 0.025 parts of triamcinolon acetonide were used and that the amounts of the stearic acid were changed to 0, 0.1, 0.3, and 0.5 parts.

- 5 The residual percentages of the triamcinolon acetonide were determined after the capsuled preparations were allowed to stand at a temperature of 60°C for one month.

The results were as shown in Table 6.

Table 6

Content of stearic acid (%)	0	0.1	0.3	0.5
Residual percentage of triamcinolone acetonide (%)	81.0	92.6	97.8	100.0

Example 12

- 20 A 100 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 1550 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$ and having a particle size such that 90% or more of the powder particles had a particle diameter of 37 to 149 μm was used.

- 25 A 10 part amount part of the hydroxypropyl cellulose and 0.17 parts of beclomethasone dipropionate were thoroughly mixed in a mixer and, then, the remaining hydroxypropyl cellulose was gradually added thereto. The resultant mixture was thoroughly mixed. Thereafter,
30 0.5 parts of stearic acid was added and thoroughly mixed.

- Thus, a powdery composition for the application to the nasal cavity containing beclomethasone dipropionate uniformly dispersed therein was prepared. The powdery composition was filled in #2 hard gelatin capsules in
35 an amount of 30 mg each to prepare a powdery preparation in a unit dosage form for the application to the nasal cavity.

Example 13

A 100 part amount of methyl cellulose having a viscosity of a 2% aqueous solution thereof of 1335 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$ was used.

5 A 10 part amount of the methyl cellulose and 0.17 parts of beclomethasone dipropionate were thoroughly mixed in a mortar and, then, the remaining hydroxypropyl cellulose was gradually added thereto. The resultant mixture was thoroughly mixed. Thereafter, 0.5 parts of
10 stearic acid was added and thoroughly mixed.

Thus, a powdery composition for the application to the nasal cavity containing beclomethasone dipropionate uniformly dispersed therein was prepared. The powdery composition was filled in #2 hard gelatin capsules in
15 an amount of 30 mg each to prepare a powdery preparation in a unit dosage form for the application to the nasal cavity.

Example 14

A powdery preparation in a unit dosage form for the
20 application to the nasal cavity was prepared in the same manner as in Example 13, except that hydroxypropylmethyl cellulose having a viscosity of a 2% aqueous solution thereof of 20 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$ was used instead of the methyl cellulose. The powdery composition was
25 filled in #2 hard gelatin capsules in an amount of 30 mg each to prepare a powdery preparation in a unit dosage form for the application to the nasal cavity.

Example 15

Powdery compositions were prepared in the same
30 manner as in Example 12, except that the amounts of stearic acid were changed to 0, 0.01, 0.1, 0.3, 0.5, and 0.7 parts. These powdery compositions were filled in #2 hard gelatin capsules in an amount of 30 mg each.

The capsules thus prepared were allowed to stand
35 at a temperature of 60°C for one month. The residual percentages of beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 7.

Table 7

Content of stearic acid (%)	0	0.1	0.3	0.5	0.7
Residual percentage of beclomethasone dipropionate (%)	77.5	87.0	93.3	96.7	98.3

10

Example 16

The same evaluation test of Example 15 was carried out, except that palmitic acid was used instead of the stearic acid.

The results were as shown in Table 8.

Table 8

Content of palmitic acid (%)	0	0.1	0.3	0.5	0.7
Residual percentage of beclomethasone dipropionate (%)	78.5	92.0	93.8	96.2	98.1

Example 17

25 Powdery compositions were prepared in the same manner as in Example 12, except that the carboxymethyl cellulose was used, instead of the stearic acid in amounts of 0, 1, 2, and 3 parts. These powdery compositions were filled in #2 hard gelatin capsules in an amount of 30 mg each.

30

The capsules thus prepared were allowed to stand at a temperature of 60°C for one month. The residual percentages of beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 9.

Table 9

Content of carboxymethyl cellulose (%)	0	1	2	3
Residual percentage of beclomethasone dipropionate (%)	82.6	92.1	93.2	95.8

Example 18

10 Powdery compositions were prepared in the same manner as in Example 13, except that the amounts of stearic acid were changed to 0, 0.01, 0.05, 0.1, and 0.5 parts. These powdery compositions were filled in #2 hard gelatin capsules in an amount of 30 mg each.

15 The capsules thus prepared were allowed to stand at a temperature of 60°C for one month. The residual percentages of beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 10.

Table 10

Content of stearic acid (%)	0	0.05	0.1	0.5
Residual percentage of beclomethasone dipropionate (%)	75.5	90.2	92.3	96.8

Example 19

30 A 90 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution of 1550 cps at 37°C \pm 0.2°C and having a particle size such that 90% or more of the particles thereof had a particle diameter of 37 to 149 μ m and 10 parts of dipotassium glycyrrhizinate were thoroughly mixed in a mixer. Then, 0.1, 0.3, 0.5,
35 and 0.7 part of stearic acid were added to the resultant mixture to prepare powdery compositions for the application to the oral cavity.

0187433

The powdery compositions prepared above were filled into #2 gelatin hard capsules in an amount of 100 mg each. The capsules thus obtained were allowed to stand at a temperature of 60°C for one month. The residual percentages of the glycyrrhizinate in the capsules were determined.

The results were as shown in Table 11.

Table 11

Content of stearic acid (%)	0	0.5	0.7
Residual percentage of glycyrrhizinate (%)	92.0	100	100

Example 20

A 100 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 1550 cps at 37°C \pm 0.2°C and having a particle size such that 90% or more of the powder particles had a particle of 37 to 149 μ m was used.

One part of the hydroxypropyl cellulose and 0.025 part of beclomethasone dipropionate were thoroughly mixed in a mortar and, then, 9 parts of hydroxypropyl cellulose were gradually added thereto. The resultant mixture was thoroughly mixed and, then, the remaining hydroxypropyl cellulose was added and thoroughly mixed in a mixer. Thereafter, 0.5 part of stearic acid was added and thoroughly mixed.

Thus, a powdery composition for the application to the oral cavity containing beclomethasone dipropionate uniformly dispersed therein was prepared. The powdery composition was filled in #2 hard gelatin capsules in an amount of 200 mg each to prepare a powdery preparation in a unit dosage form for the application to the oral cavity.

The capsules thus prepared were allowed to stand 60°C for one month and at 40°C for 6 months.

Table 12

Storage period (day)		7	14	21	30
60°C	Residual percentage of beclomethasone dipropionate (%)	100.0	99.8	98.0	97.8
Storage period (month)		1	2	3	6
40°C	Residual percentage of beclomethasone dipropionate (%)	100.0	99.0	98.7	97.5

Comparative Example 1

A 100 part amount of hydroxypropyl cellulose having a viscosity of 2% aqueous solution thereof of 1550 cps at 37°C \pm 0.2°C and having a particle size such that 90% or more of the powder particles had a particle diameter of 37 to 149 μ m was used.

One part of the hydroxypropyl cellulose and 0.025 part of beclomethasone dipropionate were thoroughly mixed in a mortar and, then, 9 parts of hydroxypropyl cellulose was gradually added thereto. The resultant mixture was thoroughly mixed and, then, the remaining hydroxypropyl cellulose were added and thoroughly mixed in a mixer. Thereafter, 0.5 part of ascorbic acid, which is different from unsaturated lower aliphatic monocarboxylic acid having six or less carbon atoms used in the present invention, was added and thoroughly mixed.

Thus, a powdery composition for the application to the oral cavity containing beclomethasone dipropionate uniformly dispersed therein was prepared. The powdery composition was filled in #2 hard gelatin capsules in an amount of 200 mg each to prepare a powdery preparation

in a unit dosage form for the application to the oral cavity.

The capsules thus prepared were allowed to stand at a temperature of 60°C for 30 days. The residual percentages of beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 13.

Table 13

Storage period (day)	7	14	21	30
Residual % of beclomethasone dipropionate	94.5	89.5	82.0	79.0

Comparative Example 2

A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Comparative Example 1, except that glutamic acid, which is different from aliphatic polycarboxylic acid used in the present invention, was used instead of the ascorbic acid.

The capsules obtained above were allowed to stand at a temperature of 60°C for 30 days. The residual percentages of the beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 14.

Table 14

Storage period (day)	7	14	21	30
Residual % of beclomethasone dipropionate	95.5	91.2	88.5	83.2

Comparative Example 3

A powd ry preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Comparative Example 1, except that
5 sodium carboxymethyl cellulose having carboxylate groups different from carboxyl groups and used instead of the ascorbic acid.

The capsules obtained above were allowed to stand at a temperature of 60°C for 30 days. The residual
10 percentages of the beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 15.

Table 15

Storage period (day)	7	14	21	30
Residual % of beclomethasone dipropionate	93.0	86.2	80.0	75.1

Comparative Example 4

A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Comparative Example 1, except that aspirin,
25 which is different from steroid or glycyrrhizic acid type anti-inflammatory agents used in the present invention, was used instead of beclomethasone dipropionate and stearic acid, which is used as a stabilizer in the present invention, was used instead of the ascorbic acid.

30 The capsules obtained above were allowed to stand at a temperature of 60°C for 30 days. The residual percentages of the aspirin in the capsules were determined.

The results were as shown in Table 16.

Table 16

Storage period (day)	7	14	21	30
Residual % of aspirin	92.2	85.5	78.2	72.0

Comparative Example 5

A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Comparative Example 1, except that azulene, which is different from steroid or glycyrrhizic acid type anti-inflammatory agents used in the present invention, was used instead of the beclomethasone dipropionate and palmitic acid, which is used as a stabilizer in the present invention, was used instead of the ascorbic acid.

The capsules obtained above were allowed to stand at a temperature of 60°C for 30 days. The residual percentages of the azulene in the capsules were determined.

The results were as shown in Table 17.

Table 17

Storage period (day)	7	14	21	30
Residual % of azulene	92.0	81.5	74.2	65.5

Comparative Example 6

A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Comparative Example 1, except that magnesium stearate, which is different from stearic acid used in the present invention, was used instead of the ascorbic acid.

The capsules obtained above were allowed to stand

at a temperature of 60°C for 30 days. The residual percentages of the beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 18.

Table 18

Storage period (day)	7	14	21	30
Residual % of beclomethasone dipropionate	94.3	90.5	85.4	80.4

Comparative Example 7

15 A 100 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 1550 cps and having a particle size distribution such that 90% or more of the powder particles had a particle diameter of 37 to 149 μ m was used.

20 A 10 part amount of the hydroxypropyl cellulose and 0.17 part of beclomethasone dipropionate were thoroughly mixed in a mixer and, then, the remaining hydroxypropyl cellulose was added thereto, followed by mixing. Thereafter, 0.5 part of ascorbic acid was added and thoroughly mixed. Thus, a powdery pharmaceutical
25 composition for the application to the nasal cavity containing beclomethasone dipropionate uniformly dispersed therein was prepared.

The powdery composition obtained above was filled in #2 hard gelatin capsules in an amount of 30 mg each
30 to prepare a powdery preparation in a unit dosage form for the application to the nasal cavity.

The residual percentages of the beclomethasone dipropionate were determined by allowing the capsules to stand at a temperature of 60°C for 30 days.

35 The results were as shown in Table 19.

Table 19

Storage period (day)	7	14	21	30
Residual % of beclomethasone dipropionate	93.6	90.1	82.2	79.1

Comparative Example 8

A powdery preparation in a unit dosage form for the application to the nasal cavity was prepared in the same manner as in Comparative Example 7, except that glutamic acid was used instead of the ascorbic acid.

The capsules obtained above were allowed to stand at a temperature of 60°C for 30 days. The residual percentages of the beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 20.

Table 20

Storage period (day)	7	14	21	30
Residual % of beclomethasone dipropionate	93.5	89.2	85.5	80.2

Comparative Example 9

A powdery preparation in a unit dosage form for the application to the nasal cavity was prepared in the same manner as in Comparative Example 7, except that aspirin was used instead of beclomethasone dipropionate and stearic acid was used instead of the ascorbic acid.

The capsules obtained above were allowed to stand at a temperature of 60°C for 30 days. The residual percentages of the aspirin in the capsules were determined.

The results were as shown in Table 21.

Table 21

Storage period (day)	7	14	21	30
Residual % of aspirin	94.2	87.2	80.5	75.1

Comparative Example 10

A powdery preparation in a unit dosage form for the application to the oral cavity was prepared in the same manner as in Comparative Example 7, except that magnesium stearate was used instead of the ascorbic acid.

The capsules obtained above were allowed to stand at a temperature of 60°C for 30 days. The residual percentages of the beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 14.

Table 22

Storage period (day)	7	14	21	30
Residual % of beclomethasone dipropionate	95.3	91.5	86.4	81.5

Example 21

A 100 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 1550 cps at 37°C \pm 0.2°C and having a particle size such that 90% or more of the powder particles had a particle diameter of 37 to 149 μ m was used.

A 10 part amount part of the hydroxypropyl cellulose and 0.17 part of beclomethasone dipropionate were thoroughly mixed in a mixer and, then, the remaining hydroxypropyl cellulose was gradually added thereto. The resultant mixture was thoroughly mixed. Thereafter, 0.5 part of stearic acid was added and thoroughly mixed. Thus, a powdery composition for the application to

the nasal cavity containing beclomethasone dipropionate uniformly dispersed therein was prepared. The powdery composition was filled in #2 hard gelatin capsules in an amount of 30 mg each to prepare a powdery preparation in a unit dosage form for the application to the nasal cavity.

The capsules thus prepared were allowed to stand at a temperature of 60°C for 30 days or at 40°C for 6 month. The residual percentages of beclomethasone dipropionate in the capsules were determined.

The results were as shown in Table 23.

Table 23

60°C	Storage period (day)	7	14	21	30
	Residual % of beclomethasone dipropionate	99.0	98.7	98.0	96.7
40°C	Storage period (month)	1	2	3	6
	Residual % of beclomethasone dipropionate	99.9	99.5	99.0	98.0

Comparative Example 11

A powdery preparation for the application to the oral cavity in the same manner as in Example 1, except that capric acid (i.e., a saturated higher aliphatic monocarboxylic acid having 10 carbon atoms) was used instead of the stearic acid.

However, the powdery preparation obtained above was not appropriate as a powdery preparation because, when the powdery preparation filled in the capsule was administered to the oral cavity by the above-explained spraying device, the odor was too strong.

Example 22

A 100 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 1550 cps at 37°C \pm 0.2°C and having a particle size such that 90%

or more of the powder particles had a particle diameter of 37 to 149 μm was used.

A 10 part amount part of the hydroxypropyl cellulose and 1.7 parts of triamcinolone acetonid were thoroughly mixed in a mixer and, then, the remaining hydroxypropyl cellulose was gradually added thereto. The resultant mixture was thoroughly mixed. Thereafter, 0.5 part of stearic acid and 0.3 part of magnesium stearate were added and thoroughly mixed. Thus, a powdery composition for the application to the nasal cavity containing triamcinolone acetonide uniformly dispersed therein was prepared.

The powdery composition obtained above was pulverized by a coarse grinding machine and, then, compressed to prepare tablets having a Monsanto hardness of 3 to 4 kg. The tablets were pulverized by the grinding machine to prepare the powder particles. The powder particles thus obtained were sieved to obtain the powder particles having a particle size of 60 to 400 meshes. The powder particles were, then, filled in #2 hard gelatin capsules in an amount of 30 mg each to prepare a powdery preparation in a unit dosage form for the application to the nasal cavity.

Example 23

A 100 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 1550 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$ and having a particle size such that 90% or more of the powder particles had a particle diameter of 37 to 149 μm was charged into a fluidized bed type granulator. Then, 50 parts of, ethanol mixture containing 0.025 part of beclomethasone dipropionate and 5 parts of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 7 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$ dissolved therein and also containing 0.5 part of stearic acid dissolved therein were sprayed from the spraying nozzle of the granulator, followed by drying. Thus, the powder particles impregnated with the solid

components in the ethanol mixture were obtained. The powder particles were sieved to obtain those having a particle size of 60 to 400 meshes. The powder particles thus obtained were filled in #2 hard gelatin capsules in an amount of 200 mg each to prepare a powdery preparation in a unit dosage form for the application to the oral cavity.

Example 24

A 80 part amount of hydroxypropyl cellulose having a viscosity of a 2% aqueous solution thereof of 1550 cps at $37^{\circ}\text{C} \pm 0.2^{\circ}\text{C}$ and having a particles size such that 90% or more of the powder particles had a particle diameter of 37 to 149 μm and 1000 parts of an ethanol suspension containing 0.2 part of the dissolved beclomethasone dipropionate and 0.4 part of the suspended stearic acid were casted on the surface of releasing paper. After drying, the casted film was obtained.

The film obtained above was ground and pulverized by a grinding machine and, then, sieved to the powder particles having a particle size of 60 to 400 meshes. The powder particles were filled in #2 hard gelatin capsules in an amount of 200 mg each to prepare a powdery preparation in a unit dosage form for the application to the oral cavity.

CLAIMS

1. A powd ry pharmaceutical composition for application to the mucosa of the oral or nasal cavity comprising:

- (a) a base selected from the group consisting of lower alkyl ethers of cellulose;
- (b) a pharmaceutically effective amount of a drug selected from the group consisting of steroid or glycyrrhizic acid type anti-inflammatory agents; and
- (c) a stabilizer for the drug composed of at least one low irritant solid organic acid selected from the group consisting of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, unsaturated lower aliphatic monocarboxylic acids having six or less carbon atoms, and cellulose derivatives having carboxyl groups.

2. A powdery pharmaceutical composition as claimed in claim 1, wherein said stabilizer is contained in an amount of 0.05% to 5% by weight in the composition.

3. A powdery pharmaceutical composition as claimed in claim 1, wherein at least about 90% by weight of the stabilizer particles has an effective particle diameter of about 75 μ m or less.

4. A powdery pharmaceutical composition for application to the mucosa of the oral cavity as claimed in claim 1, wherein said stabilizer is at least one member selected from the group consisting of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, and cellulose derivatives having carboxyl groups.

5. A powdery pharmaceutical composition for application to the mucosa of the nasal cavity as claimed in claim 1, wherein said stabilizer is a saturated higher aliphatic monocarboxylic acid having 12 or more

carbon atoms or a cellulose derivative having carboxyl groups.

6. A powdery pharmaceutical composition as claimed in claim 1, wherein said saturated higher
5 aliphatic monocarboxylic acid is a saturated higher aliphatic hydrocarbon monocarboxylic acid having 12 to 24 carbon atoms.

7. A powdery pharmaceutical composition as claimed in claim 6, wherein said saturated higher
10 aliphatic hydrocarbon monocarboxylic acid is stearic acid, palmitic acid, lauric acid, or myristic acid.

8. A powdery pharmaceutical composition as claimed in claim 1, wherein said aliphatic polycarboxylic acid is an aliphatic hydrocarbon dicarboxylic acid
15 having 4 to 6.

9. A powdery pharmaceutical composition as claimed in claim 8, wherein said aliphatic hydrocarbon dicarboxylic acid is succinic acid, fumaric acid, or maleic acid.

20 10. A powdery pharmaceutical composition as claimed in claim 1, wherein said hydroxy aliphatic polycarboxylic acid is a hydroxy aliphatic hydrocarbon di- or tricarboxylic acid having 4 to 8 carbon atoms.

25 11. A powdery pharmaceutical composition as claimed in claim 10, wherein said hydroxy aliphatic hydrocarbon di- or tricarboxylic acid is tartaric acid or citric acid.

12. A powdery pharmaceutical composition as claimed in claim 1, wherein said unsaturated lower
30 aliphatic monocarboxylic acid is an unsaturated lower aliphatic hydrocarbon monocarboxylic acid having 4 to 6 carbon atoms.

13. A powdery pharmaceutical composition as claimed in claim 12, wherein said unsaturated lower aliphatic
35 hydrocarbon monocarboxylic acid is sorbic acid.

14. A powdery pharmaceutical composition as claimed in claim 1, wherein said aromatic carboxylic

acid is benzoic acid or its derivatives.

15. A powdery pharmaceutical composition as claimed in claim 1, wherein said aromatic carboxylic acid is benzoic acid, parahydroxybenzoic acid, or
5 salicylic acid.

16. A powdery pharmaceutical composition as claimed in claim 1, wherein said cellulose derivative having carboxyl group is carboxymethyl cellulose or carboxymethylethyl cellulose.

10 17. A powdery pharmaceutical composition as claimed in claim 1, wherein said steroid type anti-inflammatory agent is beclomethasone dipropionate, betamethasone, betamethasone valerate, triamcinolone, triamcinolone acetonide, dexamethasone, fluocinolone
15 acetonide, fluocinonide, flumethasone, hydrocortisone, prednisolone, or prednisone.

18. A powdery pharmaceutical composition as claimed in claim 1, wherein said glycyrrhizic acid type anti-inflammatory agent is glycyrrhizic acid, disodium
20 glycyrrhizinate, trisodium glycyrrhizinate, monopotassium glycyrrhizinate, dipotassium glycyrrhizinate or monammonium glycyrrhizinate.

19. A powdery pharmaceutical composition as claimed in claim 1, wherein the moisture content of the
25 lower alkyl ether of cellulose is within the range of from 3% to 9% by weight.

20. A powdery pharmaceutical composition as claimed in claim 1, wherein said lower alkyl ether of cellulose is hydroxypropyl cellulose, methyl cellulose,
30 or hydroxypropyl methyl cellulose.

21. A powdery pharmaceutical composition as claimed in claim 1, wherein at least about 90% by weight of the powder particles of the pharmaceutical composition has an effective particle diameter of about 20 to 250 μ m.

35 22. A powdery pharmaceutical preparation in unit dosage form for application to the mucosa of the oral or nasal cavity comprising the powdery pharmaceutical

composition of claim 1.

23. A powdery pharmaceutical preparation as claimed in claim 22, wherein said pharmaceutical composition is filled in a capsule.

5 24. A method for preparing a powdery pharmaceutical composition for application to the mucosa of the oral or nasal cavity, comprising the step of mechanically mixing
10 (a) base selected from the group consisting of lower alkyl ethers of cellulose; (b) a pharmaceutically effective amount of a drug selected from the group consisting of steroid or glycyrrhizic acid type anti-inflammatory agents; and (c) at least one low irritant solid organic acid selected from the group consisting of saturated higher aliphatic monocarboxylic acids having
15 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, unsaturated lower aliphatic monocarboxylic acids having six or less carbon atoms, and cellulose derivatives having carboxyl groups.

20 25. A method as claimed in claim 24, wherein the mixed composition is compressed under pressure and, then, the compressed mixture is pulverized.

26. A method for preparing a powdery pharmaceutical composition for application to the mucosa of the oral or
25 nasal cavity, comprising the steps of:

impregnating the lower alkyl ethers of cellulose with a solution or dispersion, in an organic solvent, of a pharmaceutically effective amount of a drug selected from the group consisting of steroid or
30 glycyrrhizic acid type anti-inflammatory agents, and at least one low irritant solid organic acid selected from the group consisting of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic
35 polycarboxylic acids, aromatic carboxylic acids, unsaturated lower aliphatic monocarboxylic acids having six or less carbon atoms, and cellulose derivatives

having carboxyl group, and

drying the impregnated product.

27. A method for preparing a powdery pharmaceutical composition for application to the mucosa of the oral or nasal cavity comprising the steps of:

dissolving or dispersing, in an organic solvent, water, or the mixture thereof (a) the lower alkyl ethers of cellulose, (b) a pharmaceutically effective amount of a drug selected from the group consisting of steroid or glycyrrhizic acid type anti-inflammatory agents, and (c) at least one low irritant solid organic acid selected from the group consisting of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, unsaturated lower aliphatic monocarboxylic acids having six or less carbon atoms, and cellulose derivatives having carboxyl groups;

evaporating the organic solvent, water, or the mixture thereof from the resultant solution or dispersion; and

pulverizing the evaporated product.

28. A method for stabilizing a powdery pharmaceutical composition containing a steroid or glycyrrhizic acid type anti-inflammatory agent for application to the mucosa of the oral or nasal cavity, comprising incorporating at least one low irritant solid organic acid selected from the group consisting of saturated higher aliphatic monocarboxylic acids having 12 or more carbon atoms, aliphatic polycarboxylic acids, hydroxy aliphatic polycarboxylic acids, aromatic carboxylic acids, unsaturated lower aliphatic monocarboxylic acids having six or less carbon atoms, and cellulose derivatives having carboxyl groups into a mixture of lower alkyl ethers of cellulose and a pharmaceutically effective amount of a drug selected from the group consisting of steroid or glycyrrhizic acid type anti-inflammatory agents.



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
D, Y	EP-A-0 023 359 (TEIJIN LTD.) * Page 36, lines 1-9; claim 1 * & US - A - 4 294 829 ---	1-28	A 61 K 9/06 A 61 K 47/00
Y	US-A-3 276 959 (LUDWIG RITTER) * Column 5, line 54 - column 6, line 12 * ---	1-28	
A	US-A-4 289 764 (HYMAN YARROW) * Column 4, lines 26-40; claim 1 * -----	1-28	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 11-09-1985	Examiner BRINKMANN C.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	